

# Antinociceptive Activities with the Possible Mechanisms of Action of Hydroethanol Leaf Extract of *Eclipta prostrata* Hassk (Astraceae) in Mice



Esther Oluwatoyin Agbaje,\* Odunayo New Year Ashimolowo

## ABSTRACT

*Eclipta prostrata* belongs to the family of plants known as Astraceae. It is a species of plant in the sunflower family, popularly known as "false daisy" in English. The whole plant contains the alkaloids nicotine and ecliptine; and used locally for treating bruises, cuts, wounds, liver disorders and pain, among others. This study investigated the analgesic activities of hydroethanolic leaf extract of *Eclipta prostrata* in rodents using standard laboratory protocols. Doses of 50, 100, and 200 mg/kg were separately administered via oral route to the animals used in the various models - acetic acid-induced mouse writhing test, formalin - induced pain, hot plate-induced pain, and tail clip test. The effect of the extract was comparable to that produced by peripheral analgesics like aspirin and centrally acting analgesics like morphine used as positive

control in the various models employed. The mechanism of action was elucidated through pre-treatment with naloxone (1 mg/kg), yohimbine (1mg/kg), ondasetron (0.2 mg/kg), atropine (2 mg/kg), glibenclamide (2 mg/kg).

The extract activity was significant in all the models employed, however, it was more intense via the peripheral than the central pathway. Glibenclamide and atropine antagonized the activity of the decoction, which suggests that the herbal drug's antinociceptive effect is mediated via potassium ion channel opening and cholinergic receptor.

Phytochemical analysis of the leaf extract indicated the presence of flavonoids, phenols, alkaloids, phlobatanin, steroids, cardiac glycoside, and which probably contributed to the analgesic activity of the extract.

**Keywords:** antinociception, pain, medicinal plant, extract, *Eclipta prostrata*, mice

\*Correspondence to:

Prof. Mrs. E.O. Agbaje, Department of Pharmacology, Therapeutics, & Toxicology, College of Medicine, University of Lagos, Lagos, Nigeria  
[estheragbaje01@gmail.com](mailto:estheragbaje01@gmail.com)

**Cite This Article:** Agbaje, E.O., Ashimolowo, O.N.Y. 2020. Antinociceptive Activities with the Possible Mechanisms of Action of Hydroethanol Leaf Extract of *Eclipta prostrata* Hassk (Astraceae) in Mice. *Discovery Phytomedicine* 7(4): 164-170. DOI: [10.15562/phytomedicine.2020.143](https://doi.org/10.15562/phytomedicine.2020.143)

## INTRODUCTION

Pain, in its various types, is an unpleasant experience caused by intense or damaging stimuli. It is primarily protective in nature and can act as a sensorial modality to indicate the presence of tissues injury,<sup>1</sup> with its attendant negative effects on daily routine. The different exploits of pain relief has resulted in series of harmful physiological effects, including inadequate sleep, exhaustion, disorientation, anxiety, tachycardia, increased myocardial oxygen demand, immunosuppression, and increased catabolism.<sup>2</sup>

Others are respiratory depression, constipation, kidney damage, physical dependence as well as gastrointestinal irritation.<sup>3</sup> To further worsen the situation, available analgesics relieve pain as a symptom without affecting its cause,<sup>4</sup> therefore, search for new analgesic drugs with promising pharmacological actions has become an urgent need.

The use of medicinal plants as analgesic drugs in folk medicine is a common practice in developing countries; although in most cases the active principles of the plants are unknown. According to the World Health Organization over 75% of the world's population still depends on plant-derived

medicines, usually obtained from traditional healers, for its basic health-care needs. Worldwide, about 85% of primary health care medications involve the use of natural products.<sup>5</sup> Man has always discovered the great contribution of herbs in the management of almost every pharmacological disorder.<sup>6</sup>

*E. prostrata* commonly known as false daisy, (*Arojoku*; Yoruba, and *Apa*; Igbo) originated from Asia, but is widely distributed throughout the tropics and warm temperate zones. It is a much-branched, variable, prostrate, ascending or erect, rough-hairy annual herb, up to 90 cm tall with slender, reddish stems covered with short, stiff hairs, rooting at the lower nodes. Flowers throughout the year; a quantitative short-day plant, and an abundant seeder.

The various Sanskrit names of *Eclipta prostrata* are literally translated as the 'King of Hair', clearly referring to its traditional reputation in Ayurveda as an herb.<sup>7</sup> A study in rats showed that petroleum ether extracts of *E. prostrata* decreased the amount of time it took for hair to begin re-growing and to fully re-grow in shaved albino rats, thus result of treatment with *E. prostrata* was better than positive control.

Department of Pharmacology, Therapeutics, & Toxicology, College of Medicine, University of Lagos, Lagos, Nigeria

*Eclipta* species (Astraceae) have been reported useful for treating liver diseases, microbial infections, pain, constipation, respiratory diseases, urinary infections, eye problems, cardiovascular diseases, and cancer, among other disease conditions.<sup>8</sup>

The present study explored possible antinociceptive activity of *E. prostrata* and the possible pathway of producing its effect.

## METHODS AND METHODS

### Plant Material

Fresh leaves of *Eclipta prostrata* were collected from swampy area at Idi-Araba, Lagos State of Nigeria. The plant was identified and authenticated by Mr. T.K. Odewo of the Department of Botany and Microbiology, University of Lagos, Lagos, Nigeria, and given a voucher number LUH6535.

### Experimental Animals

Healthy albino mice (20-25 g) of both sexes obtained from the Laboratory Animal Centre of the College of Medicine of the University of Lagos, Idi-Araba, Lagos, Nigeria were used. The animals were maintained under standard environmental condition and were fed the standard laboratory diet, purchased from the Nigerian Institute of Medical Research (NIMR), Yaba, Lagos, alongside with clean drinking water. The animals were kept in clean plastic cages, while they acclimatized for a period of one week and were fasted for 12 hours prior to experimentation. All experimental procedures were carried out in compliance with the World Organisation for Animal Health.<sup>9</sup>

### Drugs and Chemicals

Formalin (Griffin and George, Leics, England), morphine (Martindale Pharma®, Essex, United Kingdom), acetylsalicylic acid (Reckitt Benckister Parkistan Ltd, Karachi), and acetic acid (Sigma, Chemical Company, England).

### Preparation of Plant Extract

Fresh leaves of *Eclipta prostrata* collected were rinsed under a running tap, and thereafter air dried and hand-shredded into coarse powder. A known weight, 130 g was macerated in 2 L hydroethanol (70% ethanol and 30% distilled water) for three days, and thereafter first filtered using clean white handkerchief to remove the shaft, and afterwards through a 25 mm Whatman filter paper. The filtrate was separated into different beakers and evaporated to dryness in an oven (40°C).

### Acute Toxicity Test

Groups of mice of both sexes fasted for 12 hours prior to the test were administered the extract

orally at a dose of 5000 mg/kg, while the control group receive 10 mg/kg of 1% tween 20. Doses of 10, 100, 200, 400, 800 and 1000 mg/kg extracts were administered intra-peritoneally to different groups of three animals each to determine the LD<sub>50</sub> of the plant extract. The control group was given 1% tween 20 (10 ml/kg). All animals were closely observed for toxic symptoms for the first 2 hours and mortality within 24 hour post administration was recorded. Surviving animals were observed for further 14 days post treatment for any signs of delayed toxicity. The LD<sub>50</sub> was estimated by the log dose-probit analysis method.<sup>10</sup>

## ANTINOCICEPTIVE STUDIES

### Mouse Writhing Test

The mouse writhing test was based on the method.<sup>11</sup> Albino mice of either sex weighing between 20 - 25 g were divided into six groups (n = 5). The first four received doses of extract 50, 100, 200, and 400 mg/kg, selected based on the LD<sub>50</sub>, while the remaining two groups received 1% tween 20 and 100 mg/kg aspirin respectively, one hour before intraperitoneal injection of 0.1 ml of 0.6% v/v acetic acid solution per mouse. The number of writhes/ stretches was counted for 30 minutes.

$$\% \text{ Inhibition} = \frac{\text{No of writhes control} - \text{No of writhes test}}{\text{No of writhes control}} \times 100$$

### Formalin-induced Paw Licking

Overnight-fasted mice were divided into five groups of five animals each and were treated as follows: group 1 (1% tween20 10 ml/kg), group 2 (morphine 10 mg/kg), group 3 (50 mg/kg Extract), group 4(100 mg/kg Extract) and group 5 (200 mg/kg extract). All treatments were given orally. One hour later, the mice were injected with 0.02 ml of 1% formalin subcutaneously into the right hind paw of the mice. The time (in seconds) spent in licking and biting responses of the injected paw was taken as an indicator of pain response. Response was measured for 5 minutes after formalin injection (first phase) and 15- 30 minutes after formalin injection (second phase).<sup>12</sup>

$$\% \text{ Inhibition} = \frac{\text{Reaction time control} - \text{Reaction time test}}{\text{No of writhes control}} \times 100$$

### Seconds of paw licking control

#### Hot Plate Test

The pre-reaction time to the thermal stimulus was recorded as the time interval from introduction of the animals to the plate maintained at 55± 0.5°C until the first lick of the limbs or the first jump of

the animals. The animals were initially screened and those that responded to heat stimulus within the cut-off time were randomly batched for study.

Each group was administered via oral route, the test agents as follows: group 1 (10 ml/kg 1% tween 20), group 2 (10 mg/kg morphine), group 3 (50 mg/kg extract), group 4 (100 mg/kg extract) and group 5 (200 mg/kg extract). The subsequent reactions of the animals were recorded at 30 minutes intervals for 150 minutes following the administration of all treatments.<sup>11</sup>

$$\% \text{ Inhibition} = \frac{(\text{Post-treatment Latency}) - (\text{Pre-treatment Latency})}{(\text{Cut-off Time} - \text{Pre-treatment Latency})} \times 100$$

### Haffner's Tail Clip Test

Mice that were used in this experiment were screened initially by applying a metal artery clip to the root of the tail to induce pain and animal that failed to attempt to dislodge the clip in 10 seconds were discarded. Eligible mice were divided and treated as before, using a post-treatment cut-off time of or reaction time of 30 s.<sup>11</sup> Reaction time of each mouse was determined 60 minutes post treatment for oral administration.

$$\% \text{ Inhibition} = \frac{(\text{Post-treatment Latency}) - (\text{Pre-treatment Latency})}{(\text{cut-off time} - \text{pre-treatment latency})} \times 100$$

### Elucidation of Mechanism of Antinociceptive Activity of *E. prostrata*

Randomly batched animals were respectively pre-treated with naloxone (1 mg/kg), yohimbine (1 mg/kg), ondasetron (0.2 mg/kg), atropine (2 mg/kg), glibenclamide (2 mg/kg).

Fifteen minutes post treatment, the test drug at 200 mg/kg was administered to the group of animals. Then, one hour post treatment, 0.6% v/v

acetic acid solution was injected intraperitoneally into each mouse. The number of writhes/ stretches was counted as before.<sup>13</sup>

### Data analysis and interpretation

Data was presented as mean  $\pm$  standard error of mean (SEM). Test of statistical significance was carried out using a one-way and two-way ANOVA. P value less than 0.05 ( $p < 0.05$ ) was considered statistically significant.

### Phytochemical Screening

Phytochemical screening of the crude leaf extract was carried out using standard procedures and tests to identify the phytoconstituents both qualitatively<sup>14</sup> and quantitatively. Total phenols,<sup>15,16</sup> alkaloids,<sup>17</sup> tannins,<sup>18</sup> and flavonoids.<sup>19</sup>

## RESULTS

### Physicochemical Properties of Extract

The extract was green in colour, sticky and with a pungent smell. It was readily soluble in 1% tween 20; pH 7.0.

### Acute Toxicity

Doses of 10 and 100 mg/kg produced zero percent mortality, while 200 mg/kg recorded 33.3%. Higher doses caused 100 percent deaths in the treated animals. LD<sub>50</sub> was extrapolated as 331 mg/kg. The hydroethanolic leaves extract of *Eclipta prostrata* was observed to cause moderate behavioural and sensory changes such as tachypnea, sedation, abdominal writhes, and itching.

### Acetic Acid-induced Writhing


The aqueous extract of *Eclipta prostrata* (50, 100, 200 mg/kg,) produced a significant dose-dependent ( $p < 0.05$ ) reduction in the number of writhes, when compared with the control (Figure 2). Percentage inhibition by 200 mg/kg, 100 mg/kg, 50 mg/kg of extract were 64.28%, 58.03%, 42.85% respectively, with ASA recording 71.96%.

### Formalin Test


*Eclipta prostrata* at all dose levels produced a significant dose-dependent ( $p < 0.05$ ) effect in both the phases. The highest percentage inhibition of phase one was 52.53% observed with 200 mg/kg, while 100 mg/kg produced 81.70% of the second phase. The standard drug, morphine produced an inhibition of 100% and 100% in the first and second phase respectively.

### Hot Plate Test

The extract of *Eclipta prostrata* produced a dose-dependent, but very mild action compared with the control (Table 1) and none of the dose levels



**Figure 1** LD<sub>50</sub> of *Eclipta prostrata*



**Figure 2** Effect of *E. prostrata* on acetic acid induced mouse writhing test. Data represented as mean  $\pm$  S.E.M. (n= 5). <sup>c</sup> $p < 0.05$  statistically significant compared to control (one way ANOVA followed by Dunnett's multiple comparison test)

of the extract showed any comparable effect with morphine, the standard drug used.

### Tail Clip Test

Use of the metal artery clip to the tail of animals in the control group showed reactions towards clip removal with the post-treatment latency being  $10.1 \pm 2.5$ s,  $8.51 \pm 4.38$ s,  $4.00 \pm 2.04$ s and  $8.12 \pm 5.54$ s measured at 60 minutes, 90 minutes, 120 minutes and 150 minutes respectively with a pre-treatment latency of  $2.03 \pm 1.01$ s. The percentage inhibition at 50 mg/kg are 7.54%, 17.80%, 17.09%, 13.83%. For 100 mg/kg, 2.47%, 6.83%, 13.98%, 2.75%. For 200 mg/kg, 3.9%, 14.59%, 16.3%, 28.41%. Morphine 10 mg/kg produced a peak inhibition (100%) at 90 minutes post treatment. (Table 2).

**Table 1** Effect of *E. prostrata* on Thermal Nociception

Pretreatment	Reaction time (secs)					
	Treatments	Dose (mg/kg)	60 minutes	90 minutes	120 minutes	150 minutes
Control (TW 20)	10 ml/kg	2.08 $\pm$ 0.58	1.94 $\pm$ 0.31	1.89 $\pm$ 0.34	1.74 $\pm$ 0.33	1.29 $\pm$ 0.16
<i>Eclipta prostrata</i>	50	1.54 $\pm$ 0.17	5.00 $\pm$ 0.70 <sup>ab</sup>	4.00 $\pm$ 0.31 <sup>b</sup>	2.60 $\pm$ 0.40 <sup>b</sup>	5.40 $\pm$ 0.24 <sup>ab</sup>
		Inhibition%	12.15	8.64	3.72	13.56
	100	1.58 $\pm$ 0.11	3.80 $\pm$ 0.37 <sup>b</sup>	3.40 $\pm$ 0.24 <sup>b</sup>	4.40 $\pm$ 0.81 <sup>b</sup>	5.80 $\pm$ 0.37 <sup>ab</sup>
		Inhibition%	7.81	6.40	9.92	14.84
	200	1.80 $\pm$ 0.20	3.60 $\pm$ 0.24 <sup>b</sup>	4.40 $\pm$ 0.67 <sup>b</sup>	6.80 $\pm$ 0.73 <sup>ab</sup>	3.00 $\pm$ 0.31 <sup>b</sup>
		Inhibition%	6.38	9.21	17.73	6.91
Morphine	10	3.83 $\pm$ 0.39	30.00 $\pm$ 0.00 <sup>a</sup>	25.95 $\pm$ 1.69 <sup>a</sup>	22.50 $\pm$ 1.21 <sup>a</sup>	17.51 $\pm$ 2.01 <sup>a</sup>
		Inhibition%	100	84.54	71.29	52.30

TW 20: 1% tween 20

$\pm$ S.E M: Standard Error of Mean

<sup>a</sup> $p < 0.05$  statistically significant compared to control, <sup>b</sup> $p < 0.01$  statistically significant compared to morphine (2 Way ANOVA followed by Turkey's multiple comparison tests).

**Table 2** Effect of Hydroethanolic Leaf Extract of *E. prostrata* on Tail Clip Test

Treatments	Dose (mg/kg)	Post treatments Reaction time (secs)				
		60 minutes	90 minutes	120 minutes	150 minutes	
Control (TW 20)	10 ml/kg	2.30 $\pm$ 1.01	10.10 $\pm$ 2.55	8.51 $\pm$ 4.38	4.00 $\pm$ 2.04	8.12 $\pm$ 5.54
<i>Eclipta prostrata</i>	50	1.47 $\pm$ 6.08	3.62 $\pm$ 1.56 <sup>b</sup>	6.55 $\pm$ 4.11 <sup>b</sup>	6.35 $\pm$ 2.04 <sup>b</sup>	5.42 $\pm$ 2.67 <sup>b</sup>
		Inhibition%	7.54	17.80	17.09	13.83
	100	1.51 $\pm$ 0.24	2.22 $\pm$ 0.36 <sup>b</sup>	3.44 $\pm$ 1.78 <sup>b</sup>	5.50 $\pm$ 1.79 <sup>b</sup>	2.30 $\pm$ 0.51 <sup>b</sup>
		Inhibition%	2.47	6.83	13.98	2.75
	200	1.44 $\pm$ 0.05	2.57 $\pm$ 0.15 <sup>b</sup>	5.61 $\pm$ 2.19 <sup>b</sup>	6.10 $\pm$ 1.96 <sup>b</sup>	5.69 $\pm$ 1.94 <sup>b</sup>
		Inhibition%	3.9	14.59	16.30	28.41
Morphine	10	2.21 $\pm$ 0.05	27.08 $\pm$ 1.64 <sup>a</sup>	30.00 $\pm$ 0.00 <sup>a</sup>	26.74 $\pm$ 1.36 <sup>a</sup>	23.12 $\pm$ 1.31 <sup>a</sup>
		Inhibition%	88.12	100	88.26	75.24

TW 20: 1% tween 20

$\pm$ S.E M: Standard Error of Mean

<sup>a</sup> $p < 0.05$  statistically significant compared to control, <sup>b</sup> $p < 0.01$  statistically significant compared to morphine (2 Way ANOVA followed by Turkey's multiple comparison tests).


**Table 4** Phytochemical Analysis Hydroethanolic Leaf Extract of *E. prostrata*

Chemical constituent	Inference
Phenol	+
Tannin	-
Terpernoid	-
Steroid	+
Alkaloid	+
Anthraquinone	-
Cardiac glycoside	+
Phlobatanin	+
Flavonoid	+
Saponin	-

+ = Present  
- = Absent

**Table 5** Quantitative Screening of *E. prostrata*

Chemical Compounds	Quantity (mg/100g)
Phenol	4.53
Flavonoid	11.76
Alkaloid	25.40



**Figure 3** Effect of *E. prostrata* on formalin test  
Data represented as mean  $\pm$  S.E.M. (n= 5).  $^*p < 0.05$  statistically significant compared to control (one way ANOVA followed by Dunnett's multiple comparison test)

### Mechanism of Action

The effect of the extract was reversed by glibenclamide and atropine. There was a significant increase in number of writhings when compared with the initial number recorded with the extract alone. Naloxone, yohimbine, and ondasetron did not affect the antinociceptive activity of the plant extract.

### DISCUSSION

Pain is the most common symptom of a disease; a protective mechanism to which the body

responds to harmful stimulus. It is a subjective sensory and emotional experience, connected to the stimulus that it invokes, and is also based on the observation of psychological interpretation of the phenomena taking place. The feeling of pain (nociceptive) can be caused by irritation of pain receptors, which can be found in the skin, joints, and on many internal organs. The cause of pain may also be damage to the nervous system (neurogenic), both the peripheral nerves, brain and spinal cord. Pain can also occur without damage to tissues, although the patient refers to it (psychogenic pain).<sup>20</sup>

Antinociceptive activity of *E. prostrata* hydroethanol leaf extract was investigated by peripheral/non-narcotic model of nociception such as acetic acid writhing test and central/narcotic models like hot plate and tail-flick tests. The mouse writhing test is useful for the evaluation of mild analgesic and peripherally acting drugs like aspirin (ASA), which have been reported to exhibit analgesic activity in the writhing test only. Writhings induced by acetic acid involves stimulation of the peripheral receptor system an prolonged irritation of the peritoneal cavity by acetic acid results in increase in the levels of peritoneal fluid. The analgesic effect demonstrated by hydroethanolic leaf extract of *Eclipta prostrata* in the mouse writhing test in this study suggests a peripheral mechanism of action involving direct action on nociceptors, which could possibly be via direct inhibition of prostaglandin action or an indirect action mediated through inhibition of prostaglandin synthesis by inhibiting cyclo-oxygenase (COX) activity. The effectiveness of the extract in this test indicates its potential usefulness in the treatment of acute pain as the abdominal constriction test is said to be indicative of acute pain.

The formalin test produces distinct biphasic nociceptive response; the first transient phase (early phase 0-5minutes) is caused by the direct effect of formalin on sensory C-fibers, and the second prolonged phase (late phase 15-30 minutes) is associated with the development of the injury-induced spinal sensation, responsible for facilitated pain processing; a central sensitization of the dorsal horn neuron occurred during inflammation pain.<sup>21</sup> Injection of formalin causes immediate and intense increase in the spontaneous activity of afferent C fibers a behaviour indicative of pain,<sup>22</sup> biting and licking of the injected paw as observed in this study. The formalin test can also be used to indicate the ability of analgesic agents to treat chronic pain due to inflammation. Results of the present model showed that the hydroethanolic leaf extract of *Eclipta prostrata* inhibit both the early and the late phases of formalin- induced

pain, thus suggesting its central and peripheral anti-nociceptive actions. However, from results obtained there was greater inhibition at the late phase than the early phase and this effect was comparable to that produced by the standard drug (morphine). In agreement with this, morphine was effective in both phases of the formalin test. Data from the formalin test suggests that the hydroethanol leaf extract of *Eclipta prostrata* acts through peripheral mechanism as shown in the mouse writhing test and also via central mechanism because drugs which act mainly centrally, example is morphine which is a narcotic analgesic inhibit both early and late phase of the formalin test<sup>14</sup> as demonstrated by the extract.

The investigation of the antinociceptive activity of the hydroethanolic leaf extract of *Eclipta prostrata* using the hot plate and tail clip models revealed an involvement of both  $\mu$ -opioid and  $\kappa$ -opioid receptors, since the effect was significant, although the recorded activity was mild.

Cardiac glycosides and flavonoids were among the phytoconstituents found in the plant extract. It is projected that one or a combination of the phytochemical constituents may be responsible for the analgesic activity observed with extract. Alkaloids which generally have wide pharmacological actions are analgesics and narcotics and their ability to cross the blood-brain barrier and interact with various neurotransmitter receptors determine much of their pharmacology. Some alkaloids have also been reported to possess analgesic and anti-inflammatory activity. Several reports have shown the analgesic properties of flavonoids, tannins and other polyphenolic compounds in different experimental animal models.<sup>23</sup>

In the acute toxicity test carried out for the oral route of administration of the hydroethanolic leaf extract of *Eclipta prostrata*, no mortality was observed with the 5000 mg/kg. However, the intraperitoneal route recorded a median lethal dose ( $LD_{50}$ ) of 331.5 mg/kg.

## CONCLUSION

The study demonstrated that the hydroethanolic leaf extract of *Eclipta prostrata* possess antinociceptive activity, mediated through peripheral and central pathways, using standard *in-vivo* analgesic models, and comparing its activity with conventional analgesics. The herbal drug's activity was antagonized by glibenclamide and atropine, which suggests that its antinociceptive effect is mediated via potassium ion channel opening and cholinergic receptor.

## ACKNOWLEDGEMENT

The authors express thanks to the Department of Pharmacology, Therapeutics & Toxicology, College of Medicine, University of Lagos for providing the drugs and a conducive environment for the study.

## REFERENCES

- Bektas SG, Turan S, Karadeniz U, Ozturk B, Yavas S, Biricik D, Saydam GS, & Erdemli O, Does High Thoracic Epidural Analgesia with Levobupivacaine preserve myocardium? A Prospective Randomized Study. *BioMed Res. Int.* article ID 658678, (2015) 7 pages <http://dx.doi.org/10.1155/2015/658678>.
- Helms JE, & Barone CP, Physiology and Treatment of Pain, *Crit Care Nurse* 28(6) (2008) 38-49.
- Zendeheel M, Torabi Z, & Hassanpour S, Antinociceptive mechanisms of *Bunium persicum* essential oil in the mouse writhing test: role of opiodergic and histaminergic systems, *Veterinari Medicina* 60 (2015) 63-70.
- Tripathi KD, Essentials of Medical Pharmacology 7<sup>th</sup> ed. (Jaypee Brothers Medical Publishers (P) LTD. New Delhi, India), 2013.
- Abbasi AM, Khan MA, Ahmad M, Zafar M, Jahan S, & Sultana S, (2010). Ethnopharmacological Application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, *Pakistan. J. Ethnopharmacol.* 128(2) (2010) 322-335.
- Sofowora A, Ogunbodede E, & Onayade A, The Role and Place of Medicinal Plants in the strategies for Disease Prevention Afr J Tradit Complement Altern Med 10 (5) (2013) 210-229.
- Puri HS, Rasayana: *Ayurvedic Herbs for longevity and Rejuvenation* (Taylor and Frances, London), 2003, 80-85.
- Dheba B, Vaishnavi E, Sampathkumar P, Kannan M, Hepatoprotective and Curative effects of *Eclipta prostrata* on CCl<sub>4</sub>-induced hepatotoxicity in Albino rats. *BioSc Biotech Res Asia* 9(1) (2012) 309-314.
- World Organisation for Animal Health (OIE), (2008)
- Robertson JL, Russell RM, & Preisler HK, *Bioassays with arthropods*, 2<sup>nd</sup> ed. Boca Raton, FL: CRC Press. 2007.
- Agbaje EO, Adeneye AA, & Adeleke TI, Anti-nociception and inflammatory effects of Nigerian polyherbal tonic tea (PHT) extract in rodents. *Afr J Trad Compl and Alt Med* 5(4) (2008) 399-408.
- Bhutia YD, Vijayaraghavan R, & Pathak U, Analgesic and Anti-inflammatory activity of amifostine, DRDE-07, and their analogs, in mice. *Ind. J. Pharmacol.* 42(1) (2010) 17-20.
- Fageyinbo SM, Agbaje OE, Rotimi K, Ikumawoyi V, & Fashina AY, Anti-Nociceptive Activity and Possible Mechanisms of Action of Aqueous Root Extract of *Strophanthus Hispidus* DC (Apocynaceae). *Int J Pharm Sci Res.* 7(1) (2016) 127-135. doi: [10.13040/IJPSR.0975-8232.7\(1\).127-35](https://doi.org/10.13040/IJPSR.0975-8232.7(1).127-35).
- Sofowora A, Medicinal Plants and Traditional Medicine in Africa. 3<sup>rd</sup> edn. (Spectrum Books, Ibadan), 2008.
- Mallick EP, & Singh MB, Plant Enzymology and Histoenzymology 1st ed. (New Delhi: Kalyan Publisher), 1980, 280.
- Sankhalkar S, & Vernekar V, Quantitative and Qualitative analysis of Phenolic and flavonoid content in *Moringa oleifera* Lam and *Ocimum tenuiflorum* L, *Pharmacog Res* 8(1) (2016) 16-21.
- Harborne JB, Phytochemical Methods, (Chapman Hall, London) 1998, 60-66.
- Ejikeme JCM, Ezeonu CS, & Eboatu AN, "Determination of physical and phytochemical constituents of some tropical timbers indigenous to Niger Delta Area of Nigeria," *European Scientific Journal*, 10(18) (2014) 247-270.

19. Khan AM, Qureshi RA, Ullah F, Gilani SA, Nosheen A, Sahreen S et al., Phytochemical Analysis of selected medicinal plants of Margalla Hills and surroundings, *J Med Plants Res* 5(25) (2011) 6017-6023.
20. Swieboda P, Filip R, Prystupa A, Drozd M, Assessment of pain: types, mechanism and treatment, *Ann Agric Environ Med* Special issue 1, (2013) 2-7.
21. Ashok P, Prasanna GS, & Mathuram V, Analgesic and anti-inflammatory activity of chloroform extract of *Trichilia connatoides* (W&A) Benthien. *Ind J Pharm Sc.* 68 (2006) 231-233.
22. Zakaria ZA, Abdul Ghani ZDF, Raden Mohd Nor RNS, Gopalan HK, Sulaimon Mohd R, Mat Jais AM, Somchit MN, Kader AA, & Ripin J, Antinociceptive, anti-inflammatory, and antipyretic properties of an aqueous extract of *Dicranopteris linearis* leaves in experimental animal model *J Nat Med.* 62(2) (2008) 179-87. DOI: [10.1007/s11418-007-0224-x](https://doi.org/10.1007/s11418-007-0224-x).
23. Woode E, Amoh-Barimah AK, Abotsi WK, Ainooson GK, & Owusu G, Analgesic effects of stem bark extracts of *Trichilia monadelpha* (Thonn.) JJ De Wilde. *Ind J Pharmacol* 44(6) (2012) 765-773.



This work is licensed under a Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>