

Antisickling and antibacterial activities of *Uvariopsis congensis* Robyns & Ghesq. (Annonaceae)



Koto-te-Nyiwa Ngbolua^{1,2,*}, Lengbiye E. Moke¹, Masengo C. Ashande², Justin A. Asimonyio³,
Jean Leon K. Kambale³, Damien S.T. Tshibangu¹, Nadège K. Ngombe⁴, Ilunga Kamika⁵,
Pius T. Mpiana¹

ABSTRACT

The aim of the present study was to evaluate the chemical composition and bioactivity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract against Sickle cell disease and associated bacteria. This plant species ingested by wild chimpanzees as medicinal food could inhibit the human hemoglobin S polymerization inducing hemolysis of sickle erythrocytes (as it does by inhibiting the *Plasmodium falciparum* inducing hemolysis of infected erythrocytes in great apes). The antisickling and antibacterial activities were assessed according respectively to Emmel and micro-dilution methods. The results revealed that *Uvariopsis congensis* Robyns & Ghesq. stem bark extract contains various secondary metabolites such as alkaloids, saponins, total polyphenols ($87.942 \pm 0.536 \mu\text{g GAE/g}$), flavonoids ($0.623 \pm 0.630 \mu\text{g QE/g}$), tannins ($0.889 \pm 0.002 \mu\text{g TAE/g}$), anthocyanins ($0.150 \pm 0.017 \mu\text{g M-3GE/g}$), leuco-anthocyanins, quinones, terpenes and steroids. Ethanolic extract displayed interesting antisickling activity which was confirmed by the appearance of radius, the increase of the cell area and the decrease of its perimeter ($p < 0.05$). The antisickling activity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract is dose dependent. The minimal concentration of normalization (MCN) and the dose of extract for which 50% of the sickled erythrocytes are reversed (ED50) were $100 \mu\text{g/mL}$ and $0.6 \mu\text{g/ml}$ respectively. The minimum inhibitory concentrations (MIC) were $250 \mu\text{g/mL}$ and $500 \mu\text{g/mL}$ respectively for *Escherichia coli* ATCC 27195 and *Staphylococcus aureus* ATCC 33591. The bioactivity of this plant species could be improved by liquid-liquid extraction and chromatographic bio-guided fractionation of ethanolic extract. This study validates for the first time the *in vitro* antisickling activity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract. The animal self-medication constitutes thus an innovative method which could lead to the discovery of new drugs for the treatment of human ailments like Sickle cell disease.

Keywords: Sickle cell disease, Congo basin, great apes plant foods, zoopharmacognosy, novel drugs

Introduction

Infectious diseases are recognized to affect the behavior and reproductive fitness of all animals.¹ As response, the animal kingdom developed anti-parasitic behaviors through plant-animal co-evolution relationships in which some animal phylum utilize the chemical defenses of plants to protect themselves from their own parasites.²

In African tropical forests like Congo basin, Great apes (GAs) were reported as a good model for the understanding of malaria infection patho-physiology. Indeed, Humans and great apes (bonobos, chimpanzees, gorillas, etc.) share a common gut anatomy. Although, some diseases that cause countless deaths in humans (like malaria) are ineffective or have minor non disturbing effects in GAs. These animals adopt a self medicative behavior when they are displaying malaria symptoms by selecting specific plants for

controlling parasite infection while this one cause hemolytic anemia of human red blood cells.^{3,4}

In this regards, we recently hypothesized that these plant items could therefore protect human sickle erythrocyte against hemolysis (by inhibiting the polymerization of sickle hemoglobin and radical oxygen species formation within sickle erythrocyte) as it does for *Plasmodium falciparum* infected erythrocytes in great apes.^{5,6} So, some of such plant species occasionally ingested by GAs for their supposed medicinal properties were reported to contain natural product compounds that prevent the erythrocytes hemolysis through the inhibition of sickle erythrocyte sickling and this bioactivity was scientifically validated.⁶⁻¹¹

Uvariopsis congensis is a shrub or small tree. The flowers are solitary and borne in the axils of the leaves and on the older wood. The leaf is smaller, long by 2 to 6 cm, broad and is elliptic instead of oblong to oblanceolate and entirely glabrous. The flowers and fruits are very shortly

*Corresponding author, E-mail: jpngbolua@unikin.ac.cd; ¹Department of Biology, ²Research for Sustainable Development, ³Biodiversity monitoring centre, ⁴Faculty of Pharmaceutical Sciences, Faculty of Science University of Kinshasa, Democratic Republic of the Congo; ⁵Department of Environmental Sciences, College of Agriculture and Environmental Science, University of South Africa, UNISA, South Africa. Copyright: © 2016 The Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License.

pedunculate; the petals ovate and sub-acute (instead of rotund). The fruits are pinkish red in color, about 1.5 cm diameter and constricted between the seeds.¹²

The plant species was reported to be eaten by wild chimpanzees (*Pan troglodytes schweinfurthii*).¹³ The active compounds (acetogenins) isolated from *Uvariopsis congensis* Robyns & Ghesq. stem bark extract showed highly significant medicinal properties like antibacterial, antimalarial, and/or antileishmania.¹⁴ However, no study has yet been reported on the antisickling activity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract. The plant could be potentially non-toxic to man and could provide new sources of anti-sickling compounds.

Sickle cell disease (SCD) is a hemoglobinopathy which naturally occurring in black people. At the genetic level, SCD is characterized by a single base substitution in the gene encoding the human β -globin subunit and results in replacement of $\beta 6$ glutamic acid by valine that causes an abnormal, rigid and sickle shape in hypoxia.^{15,16}

In Democratic Republic of the Congo (DRC), surveys reported that 12% of the hospitalized children are sicklers and the annual cost of the treatment of SCD is higher than 1.000,00 USD per patient.¹⁷ It is therefore necessary to develop innovative strategy for identifying new source of antisickling agents through zoopharmacognosy.⁶⁻¹¹

The aim of this study is to evaluate the chemical composition and bioactivity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract against SCD and associated bacteria, a plant species which has been selected through the animal source of knowledge (new paradigm) in order to discover new phytochemistry for human.

Materials and Methods

Plant identification and preparation of extracts

The plant material (stem barks) used in the present study was collected in Kisangani city (Tshopo Province, Democratic Republic of the Congo) during a field work in July-August 2015 and authenticated by Justin A. Asimonyio of the Biodiversity monitoring centre, University of Kisangani. Voucher specimen No JAA05T is on deposit at the Laboratory of Molecular bioprospection (Department of Biology, Faculty of Science of the Kinshasa University).

Test extract preparation

The plant material (1 kg) was kept at room temperature (25 to 30 °C) for air drying (two weeks). The air-dried and powdered material (50 g) was extracted by repeated maceration with ethanol 90° (3×4 hrs, 1.5 L) at room temperature. After filtering the mixture, the aqueous-ethanol filtrates were pooled, dried over Na₂SO₄ and evaporated to dryness under reduced pressure using a rotary evaporator to yield crude ethanolic extract. Extract was stored at 4 °C.

Phytochemical screening

The dried and powdered plant material (10 g) was repeatedly extracted by cold percolation with 95% ethanol (EtOH) and water (100 mL × 2) for 48 hours. Chemical screening was performed on the aqueous and organic extracts to investigate the presence of alkaloids, saponins, total polyphenols, flavonoids, tannins, anthocyanins, leucoanthocyanins, quinones, terpenes and steroids according to standard protocol.¹⁸

Detection of phenols (Ferric Chloride Test)

Total phenolics contents were determined according to the Folin-Ciocalteu method with slight modifications.¹⁹ The extract (200 μ L) was mixed with 1.5 mL of Folin-Ciocalteu reagent (previously diluted 10 times with double distilled water) and allowed to stand at room temperature for 5 min. 1.5 mL of sodium bicarbonate solution (60 g/L) was added to the mixture and after incubation for 90 min at room temperature, the absorbance was measured at 725 nm using a UV-Visible spectrophotometer (GENESYS 10S). Total phenolic were quantified by calibration curve obtained from measuring the absorbance of the known concentrations of gallic acid standard solutions (10-150 μ g/mL in 80% methanol). The results were calculated as gallic acid equivalent (GAE) per one gram dry powder and reported as mean value \pm standard deviation (SD) (the standard curve equation: $Y = 0,006 \times -0,002$; $R^2 = 0,997$).

Detection of flavonoids

Total flavonoid content was measured by the aluminum chloride colorimetric method. 20 An aliquot (1 mL) of each extract was added to 10 mL volumetric flask containing 4 mL of double distilled water. Then 0.3 mL NaNO₂ 5% was added to the flask and after 5 min, 0.3 mL AlCl₃ (10%) was also added. At 6th min, 2 mL NaOH (1 M) was added and the total volume was made up

to 10 mL with double distilled water. The solution was mixed completely and the absorbance was measured versus prepared reagent blank at 510 nm. Total flavonoid content was expressed as mg quercetin equivalents (QE) per one gram dry powder. One mL of standard solution (quercetin: 5-100 µg/mL) was used to construct calibration curve (the standard curve equation: $Y = 0.009 \times + 0.006$; $R^2 = 0,999$).

Detection of tannins

To 1 ml of the extract was added 7,5 ml of distilled water and 0,5 ml of Folin-Ciocalteu reagent and 1 ml of sodium carbonate (Na₂CO₃ 35%). The absorbance was measured at the wavelength of 725 nm. The tannins content (expressed as tannic acid equivalent, TAE) was calculated using the following relation: $Y = 0,443 \times - 0,264$; $R^2 = 0,720$.²²

Detection of anthocyanins

The samples were diluted with the mixture ethanol/water/HCl conc. (70:30:1; v/v/v) and the absorbance was measured at the wavelength of 540 nm. The anthocyanins content (expressed as malvidin-3-glucoside equivalent, M-3-GE) was calculated using the following relation: Anthocyanins = $A_{540} \times (10/0.6) \times d$ (with A_{540} = maximum of absorption at 540 nm; d = dilution factor; 0.6 maximum of absorption of 10 mg/L of M-3-GE standard solution).²¹

Bioassays

In vitro antisickling bioassay

Blood samples used to assess the antisickling activity of the selected plant extracts were taken from known SCD patients attending the "Centre de Médecine Mixte et d'Anémie SS" located in Kinshasa, Democratic Republic of the Congo. None of the patients had been transfused recently with Hb AA blood and all antisickling experiments were carried out with freshly collected blood. In order to confirm their SS nature, the above-mentioned blood samples were first characterized by hemoglobin electrophoresis on cellulose acetate gel to confirm their status and were then stored at 4 °C in a refrigerator. An informed consent was obtained from all the patients participating in the study and all the research procedures have received the approval of Department of Biology Ethics Committee.

An aliquot of Hb S-blood was diluted with 150 mM phosphate buffered saline (NaH₂PO₄ 30 mM, Na₂HPO₄ 120 mM, NaCl 150 mM) and mixed with an equivalent volume of 2%

sodium metabisulfite. A drop from the mixture was spotted on a microscope slide in the presence or absence of plant extracts and covered with a cover slip. Paraffin was applied to seal the edges of the cover completely to exclude air (Hypoxia). Duplicate analyses were run for each extract. The red blood cells (RBCs) were analyzed by a computer assisted image analysis software (Motic Images 2000, version 1.3; Motic China Group Co LTD) and statistical data analysis were processed using Microcal Origin 8.5 Pro package software as previously reported.²³⁻²⁷

Determination of antibacterial activity

The activity of the plant extract was tested against *Staphylococcus aureus* (*S. aureus* ATCC 33591) and *Escherichia coli* (*E. coli* ATCC 27195) strains. The tested strains were obtained from the American Type Culture Collection (ATCC, Rockville MD, USA)

Determination of Minimum inhibitory concentration (MIC)

The Minimum inhibitory concentration (MIC) was determined by broth micro-dilution method as reported in our previously research work.^{28, 29} The inocula of used microorganisms were prepared from 24 hours old broth cultures. The absorbance was read at 600 nm and adjusted with sterile physiological solution (0.9% NaCl) to match that of a 0.5 McFarland standard solution (108 cells/mL). The prepared microbial suspension was diluted (1/100) to achieve 106 CFU/mL. Stock solutions of the plant extracts were prepared in Tween 80 (Fisher chemicals) (3 mg/300 µL) and diluted to 2.7 ml with Mueller Hinton Broth (MHB) (Conda, Madrid, Spain) to achieve a Tween 80 final concentration of 0.1%. This solution was transferred in 96-wells plates (200 µL/well) and two-fold serially diluted with MHB to give final concentrations ranging from 1000 to 3,906 µg/mL.

An aliquot (10 µL) of 106 CFU/mL overnight culture was added to wells of a sterile 96-well micro-plate titer. The positive control wells contained both MHB and bacteria suspension without plant extract while negative control wells contained MHB only. The MIC was determined as the lowest plant extract concentration at which no growth were observed after 24 hours. Resasurin (30 µL) in aqueous solution (0.01%) was used to evaluate the micro-organism viability.

Statistical analysis

The results of in vitro study are given as Mean \pm Standard Deviation obtained from three independent experiments. The results of in vivo study were expressed also as Mean \pm Standard Deviation and analyzed with Student's t-test for paired data using Origin version 8.5 Pro package software. All data were analyzed at a 95% confidence interval ($\alpha = 0.05$).

Results and Discussion

Extract yields and chemical screening

The phytochemical analyses performed on *Uvariopsis congensis* Robyns & Ghesq. stem bark extract revealed the presence of alkaloids, saponins, total polyphenols ($87.942 \pm 0.536 \mu\text{g}$

GAE/g), flavonoids ($0.623 \pm 0.630 \mu\text{g QE/g}$), tannins ($0.889 \pm 0.002 \mu\text{g TAE/g}$), anthocyanins ($0.150 \pm 0.017 \mu\text{g M-3GE/g}$), leuco-anthocyanins, quinones, terpenes and steroids. Phenolic compounds such as anthocyanins^{3, 9, 16, 23-27, 31-33} rosmarinic acid³⁴ and lunularic acid³⁵ and triterpenes like betulinic, maslinic, oleanolic³⁶ and ursolic acid³⁷ were reported to display antisickling activity *in vitro*.

Antisickling activity

The figure 1 give the morphology of untreated sickle erythrocytes (SE) and SE treated with *Uvariopsis congensis* Robyns & Ghesq. stem bark extract.




Figure 1. Morphology of untreated sickle erythrocytes (a) and SE treated with 50 $\mu\text{g/mL}$ of *Uvariopsis congensis* (b) (X500), [NaCl 0.9%; Na₂S₂O₅ 2%].

Cell parameters	Untreated SE	SE + EtOH
Radius (μm)	0.00 \pm 0.00	3.5 \pm 0.4
Perimeter (μm)	36.0 \pm 1.5	19.2 \pm 1.1
Area (μm^2)	21.4 \pm 1.3	34.6 \pm 1.2

Table 1. Average values of radius, perimeter and surface of untreated and sickle erythrocytes (SE) treated with ethanolic extract of *Uvariopsis congensis*.

Figure 1a revealed that the control contains in majority sickle-shaped erythrocytes, confirming the homozygous (SS) nature of the blood. Mixed together with ethanolic extract (Figure 1b), the majority of erythrocytes are reversed normal-shape. This indicates that *Uvariopsis congensis* Robyns & Ghesq. stem bark extract have antisickling effect. Our previous research works revealed that this activity could be due to compounds such as anthocyanins, phenolic or triterpenic acids.^{3, 9, 16, 23-27, 30-37} The treated SE demonstrated a remarkable similarity to normal red blood cells.

The table 1 gives the average values of cell parameters (radius, perimeter and area) of

untreated and sickle erythrocytes treated with the extract of *Uvariopsis congensis*.

Table 1 indicates that untreated sickle cells are not circular, so it's not possible to calculate the radius due to their sickled shape in hypoxic conditions. While, in the presence of plant extract; drepanocyte recovered normal shape.

This phenomenon is confirmed by the appearance of radius, the increase of the treated cell area and the decrease of it perimeter if compared to untreated sickle erythrocyte ($p < 0.05$). Indeed, the computer software package (Motic Images 2000 version 1.3) used in this study did not give the average radius for untreated sickle




Figure 2. Dose-dependent normalization of sickle erythrocytes treated with extract from *Uvariopsis congensis*

cells, as sickled cells of untreated blood were not circular. The average radius appeared after treatment of sickle cells with ethanolic extract of *Uvariopsis congensis* is an indication of the bioactivity of this plant species.

The normalization rate of sickled cells increases with the extract concentration and reached a maximum and constant value at 100 µg/mL (minimal concentration of normalization, MCN). This corresponds to a normalization rate of 89% with an ED50 (dose of extract for which 50% of the sickled erythrocytes are reversed) equal to 0.6 µg/mL. Thus, the antisickling activity of *Uvariopsis congensis* is dose dependent (figure 2). The extracts from *Uvariopsis congensis*, because of their bioactive secondary metabolites richness like phenolic compounds (anthocyanins), could exert the antisickling effect through various mechanisms like the inhibition of free radicals formation, the inhibition of hemoglobin polymerization, and the inhibition of erythrocyte hemolysis.

Antibacterial activity

Due to the high cost of modern therapy for SCD, a medicinal plant species displaying at the same time antisickling and antibacterial activities could be useful in the management of this disease. The antibacterial activity of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract was evaluated against *E. coli* and *S. aureus* strains.

Results revealed that the negative bacterium *E. coli* was more sensitive to plant extract (MIC=250 µg/mL) than *S. aureus* (MIC=500 µg/mL). This antibacterial activity is however weak if compared to the MIC reference value of 100 µg/mL. Indeed, a value lower or equal to 100 indicates a great antibacterial activity. The difference in the bioactivity would be due to the nature of the bacterial wall. These results

are in agreement with previous works on the antimicrobial activity of natural products of plant origin. It was reported that *S. aureus* and *E. coli* constitute both the principal bacteria responsible for septicemia and the osteomyelitis in SCD patients. To this end, *Uvariopsis congensis* is a better candidate for the development of phytomedicine with broad spectrum of action for the management of SCD. The antibacterial activity observed in this study can be improved through liquid-liquid extraction and chromatographic bioguided fractionation of the ethanolic extract. To our knowledge, it is for the first time that the antisickling activity of *Uvariopsis congensis* is reported in the literature.

Conclusions

The present study evaluated the chemical composition and the antisickling and antibacterial activities of *Uvariopsis congensis* Robyns & Ghesq. stem bark extract. The results revealed that: The stem bark of *Uvariopsis congensis* contains various secondary metabolites such as the anthocyanins, flavonoids, tannins, quinones, saponins, alkaloids, steroids, terpenoids and leuco-anthocyanins; Ethanolic extracts displayed very interesting antisickling activity in vitro (ED50= 0.6 µg/mL). These extracts could exert their antisickling effect through various mechanisms including the inhibition of free radicals formation, the inhibition of hemoglobin polymerization, and the inhibition of erythrocyte hemolysis. The minimum inhibitory concentrations (MIC) were 250 µg/mL and 500 µg/mL respectively for *E. coli* ATCC 27195 and *S. aureus* ATCC 33591.

This study provided experimental evidence that supports further development of *Uvariopsis congensis* extracts as a medicine for the management of SCD in the Congo basin. The animal self-medication constitutes thus an

innovative method which could lead to the discovery of new drugs for the treatment of human ailments.

Conflict of Interest

None

Acknowledgements

The authors thank the “Académie de Recherche et d'Enseignement supérieur (ARES), Belgium” for Research grants PAH ARES/UNIKIN 2015 offered to Koto -te- Nyiwa Ngbolua, Pius T. Mpiana and Damien S.T. Tshibangu. The authors are also indebted to The World Academy of Sciences (TWAS) (Grant No. 15-156 RG/CHE/AF/AC_G–FR3240287018) for providing financial assistance.

References

- Huffman MA. Animal origins of herbal medicine. In: “From the sources of knowledge to the medicines of the future” Jacques Fleurentin, Jean-Marie Pelt, Guy Mazars eds, Proceedings of the 4th European Congress on Ethnopharmacology, 11-13 may 2000, Metz (France), pp. 31-42.
- Huffman MA. Current evidence for self-medication in primates: a multidisciplinary perspective. *Phys. Anthropol.* 1997; 40: 171-200.
- Ngbolua KN. Evaluation de l'activité anti-drépanocytaire et antipaludique de quelques taxons végétaux de la République Démocratique du Congo et de Madagascar, Thèse de Doctorat: Université de Kinshasa, République Démocratique du Congo, 2012. DOI : 10.13140/RG.2.1.3513.3606.
- Krief S, Escalante AA, Pacheca MA, Mugisha L, André C. On the diversity of malaria parasites in African apes and the origin of *P. falciparum* from bonobos. *PLoS Pathog.* 2010; 6(2): e 1000765.
- Krief S. La pharmacopée des chimpanzés. *Pour La Sci.* 2004; 325:76–80.
- Kambale JK, Ngbolua KN, Mpiana PT, Mudogo V, Tshibangu DST, Wumba R, Mvumbi LG, Kalala LR, Mesia KG, Ekutsu E. Evaluation in vitro de l'activité antifalcémiant et effet antioxydatif des extraits d'*Uapaca heudelotii* Baill. (Euphorbiaceae). *Int. J. Biol. Chem. Sci.* 2013 ; 7(2): 523-534.
- Ngbolua KN, Bolaa BM, Mpiana PT, Ekutsu EG, Masengo AC, Tshibangu DST, Mudogo V, Tshilanda DD, Kowozogono KR. Great Apes Plant Foods as Valuable Alternative of Traditional Medicine in Congo Basin: The Case of Non-Human Primate Bonobos (*Pan paniscus*) Diet at Lomako Fauna Reserve, Democratic Republic of the Congo. *J. Adv. Bot. & Zool.* V311, 2015a. DOI: 10.15297/JABZ.V311.01.
- Ngbolua KN, Tshibangu DST, Mpiana PT, Mihigo OS, Mavakala BK, Masengo CA, Muanyishay LC. Anti-sickling and antibacterial activities of Some Extracts from *Gardenia ternifolia* subsp. *jovis-tonantis* (Welw.) Verdc. (Rubiaceae) and *Uapaca heudelotii* Baill. (Phyllanthaceae). *J. Adv. Med. & Pharm. Sci.* 2015b; 2 (1): 10-19.
- Ekutsu GE, Ngbolua KN, Bolaa MB, Mpiana PT, Ngoy BP, Masengo AC, Bongo GN. Enquête sur la pharmacopée des bonobos (*Pan paniscus*, Primates) dans un foyer endémique et Mise en évidence de l'activité anti-drépanocytaire chez un taxon végétal (*Treulia africana* Decne ex Trécul, Moraceae) testé in vitro. *Int. J. Innov. & Appl. Studies* 2016; 14 (2): 315-326.
- Ngbolua KN, Bishola TT, Mpiana PT, Mudogo V, Tshibangu DST, Ngombe KN, Ekutsu EG, Gbolo BZ, Kabena NO. Ethno-pharmacological survey, in vitro antisickling and free radical scavenging activities of *Carapa procera* DC. stem bark (Meliaceae). *Nova J. Med. & Biol. Sci.*, 2014; 2 (2): 01-14.
- Ngbolua KN, Bishola TT, Mpiana PT, Mudogo V, Tshibangu DST, Ngombe KN, Ekutsu EG, Tshilanda DD, Gbolo BZ, Mwanangombo TD, Fatiany PR, Baholy R. Ethno-botanical survey, in vitro antisickling and free radical scavenging activities of *Garcinia punctata* Oliv. (Clusiaceae). *J. Adv. Bot. & Zool.* V112, 2014. DOI: 10.15297/JABZ.V112.04.
- Lemmens RHMJ. *Uvariopsis congensis* Robyns & Ghesq. In: Lemmens RHMJ, Louppe D & Oteng-Amoako AA (Editors). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands. Consulté le 24 mai 2016.
- Krief S, Huffman MA, S'evenet MA, Hladik CM, Grellier P. Bioactive properties of plant species ingested by chimpanzees (*Pan troglodytes schweinfurthii*) in the Kibale National Park, Uganda. *Am. J. Primatol.* 2006; 68: 51-71.
- Krief S, Hladik CM, Haxaire C. Ethnomedicinal and bioactive properties of plants ingested by wild chimpanzees in Uganda. *J. of Etnopharmacol.* 2005; 101: 1-15.
- Giroto R, Begué P, Galacteros F. La drépanocytose. Editions John LIBBEY Eurotext, Paris: France, 2003.
- Mpiana PT, Ngbolua KN, Bokota MT, Kasonga TK, Atibu EK, Mudogo V. In vitro Effects of Anthocyanins Extracts From *Justicia secunda* VAHL on the Solubility of Hemoglobin S and Membrane Stability of Sickle Erythrocytes, *Blood Transfus.* 2010; 8: 248-254.
- Tshilolo L, Aissi LM, Lukusa D, Kinsiyama C, Wembonyama S, Gulbis B, Vertongen F. Neonatal screening for sickle cell anaemia in the Democratic

- Republic of Congo: experience from a pioneer project on 31204 newborns. *J. Clin. Pathol.* 2009; 62:35-38.
18. Bruneton J. *Pharmacognosie, Phytochimie des Plantes Médicinales*. 3rd Edition, Revue et Augmentée, Tec & Doc, Paris, 1999.
 19. Gutfinger T. Polyphenols in olive oils. *J. Am. Oil Chem. Soc.* 1981; 58: 966-968.
 20. Quertier-Deleu C, Gressier B, Vasseur J, Dine T, Brunet C, Luyckx MC, Cazin JC, Bailleu F, Trotin F. Phenolic compounds and antioxidant activities of buckwheat (*Fagopyrum esculentum* Moench) hulls and flour. *J. Ethnopharmacol.* 2000; 72: 35-42.
 21. Di Stefano R, Cravero MC, Gentilini N. *Meto-di per lo studio dei polifenoli dei vini*. *L'enotecnico*. 1989 ; 5: 83-90.
 22. Makkar HPS, Bluemmel M, Borowi NK, Beker K. Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods. *J. Sci. Food Agric.* 1993; 61: 161-165
 23. Mpiana PT, Mudogo V, Nyamangombe L, Kakule MK, Ngbolua KN, Atibu EK, Mbala MB, Mbongo AK, Ntumba JN. Antisickling Activity and Photodegradation Effect of Anthocyanins Extracts from *Alchornea cordifolia* (SCHUMACH & Thonn.) and *Crotalaria retusa* L. *Ann. Afr. Med.* 2009; 2 (4): 240-245.
 24. Mpiana PT, Misakabu FM, Kitadi JM, Ngbolua KN, Tshibangu DST, Lombe BK, Tsalu PV, Atibu EK, Gbolo BZ, Muanishay CL. Antisickling activity and physico-chemical stability of anthocyanin extracts from *Hypoxis angustifolia* Lam. (Hypoxidaceae) bulbs. M. Noboru (Ed.), in: *Anthocyanins: Occurrence, Structure, Biosynthesis and Health benefits Based on their Evidences of Phytochemicals in Vegetables and Fruits*, NOVA Science Publishers, Inc., New York, USA, 2014; Vol. 2, pp: 97-113.
 25. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Atibu EK, Kitwa EK, Kanangila AB, Makelele LK. *Activité antifalcémiant et thermodégradation d'une fraction d'anthocyanes extraits de Zizyphus mucronata*, *Ann. Afr. Med.* 2009; 2 (2) : 91-97.
 26. Mpiana PT, Mudogo V, Tshibangu DST, Shetonde OM, Ngbolua KN, Mbala MB. Antisickling activity of some Congolese plants, in: *Drug discovery from African flora, The 12th Symposium of the Natural Product Research Network for Eastern and Central Africa*, July 22-26, 2007; University of Makerere, Kampala, Uganda, p.45 (PS-6).
 27. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Mangwala PK, Atibu EK, Kakule MK, Makelele LK, Bokota MT. Antisickling Activity and Thermodegradation of an Anthocyanin fraction from *Ocimum basilicum* L. (LAMIACEAE). *Comp. Bio. Nat. Products*. 2010; Vol. 3. Effects, Safety & Clinical (Pt II), pp. 278-287.
 28. Ngbolua KN, Mubindukila NR, Mpiana PT, Masengo CA, Baholy R, Fatiany PR, Ekutsu GE, Gbolo BZ. In vitro Assessment of Antibacterial and Antioxidant activities of a Congolese medicinal plant species *Anthocleista schweinfurthii* Gilg (Gentianaceae). *J. Mod. Drug Discov. & Drug Deliv. Res.* 2014; V113. DOI: 10.15297/JMDDR.V113.03.
 29. Ngbolua KN, Mubindukila NR, Mpiana PT, Tshibangu DST, Masengo CA, Nzongola K, Baholy R, Fatiany PR. Phytochemical screening, Antibacterial and Antioxidant activities of *Anthocleista liebrechtsiana* Wild & T. Durand (Gentianaceae) originated from Democratic Republic of the Congo. *J. Adv. Med. & Life Sci.* V113, 2014; DOI: 10.15297/JALS.V113.04.
 30. Ngbolua KN, Mudogo V, Mpiana PT, Malekani MJ, Rafatro Herintsoa, Ratsimamanga S Urverg, Takoy L, Rakotoarimana H, Tshibangu DST. Evaluation de l'activité anti-drépanocytaire et antipaludique de quelques taxons végétaux de la République démocratique du Congo et de Madagascar. *Ethnopharmacol.* 2013; 50: 19-24.
 31. Ngbolua KN, Mpiana PT, Mwanza BF, Tshibangu DST, Tshilanda DD, Masengo AC, Mudogo V, Baholy R, Fatiany PR. Antisickling and antibacterial activities of *Garcinia punctata* Oliv. (Clusiaceae) and *Tetradenia riparia* (Hochst.) Codd (Lamiaceae) from Democratic Republic of the Congo. *J. Adv. Med. & Life Sci.* V411. DOI: 10.15297/JALS.V411.01, 2016.
 32. Mpiana PT, Ngbolua KN, Mudogo V, Tshibangu DST, Atibu EK, Tshilanda DD, Misengabu NM. Antisickle erythrocytes haemolysis properties and inhibitory effect of anthocyanins extracts of *Trema orientalis* (ULMACEAE).
 33. Mpiana PT, Kimbadi BL, Ombeni AM, Ngbolua KN, Tshibangu DST, Tshilanda DD, Misengabu NM, Muanyishay CL, Muyisa SK. In vitro inhibitory effects and anti-sickle erythrocytes haemolysis of *Dicliptera colorata* C.B. Clarke, *Euphorbia hirta* L. and *Sorghum bicolor* (L.) Moench. *Open J. Blood Dis.* 2013; 3: 43-48.
 34. Tshilanda DD, Onyamboko DVN, Babady PB, Mutwale PK, Tsalu PV, Tshibangu DST, Ngombe NK, Ngbolua KN, Mpiana PT. Chemical fingerprint and anti-sickling activity of Rosmarinic acid and Methanolic extracts from three Species of *Ocimum* from DR Congo. *J. Biosci. & Med.* 2016; 4: 59-68.
 35. Ngbolua KN, Rafatro H, Rakotoarimanana H, Mudogo V, Mpiana PT, Tshibangu DST, Tshilanda DD. In vitro anti-erythrocyte sickling effect of lunularic acid of natural origin. *Int. Blood Res. & Rev.* 2015; 4(3):1-6.
 36. Tshibangu DST, Shode FO, Koorbanally N, Mudogo V, Mpiana PT, Ngbolua KN. Antisickling triterpenoids from *Callistemon viminalis*, *Meulaleuca bracteata* var. *Revolution Gold* *Syzygium guineense* and *Syzygium cordatum*. The

- 14th NAPRECA Symposium and AAMPS Ethnoverterinary Medicine Symposium 8th-12th August 2011. International Centre For Insect Physiology and Ecology (ICIPE): Kasarani, Nairobi, Kenya, pp.296-300 (YS 27).
37. Tshilanda DD, Onyamboko DNV, Babady PB, Ngbolua KN, Tshibangu DST, Dibwe EF, Mpiana PT. Anti-sickling Activity of Ursolic Acid Isolated from the Leaves of *Ocimum gratissimum* L. (Lamiaceae). *Nat. Prod. Bioprospect.* 2015; 5: 215-221.
38. Ngbolua KN, Mpiana PT, Akoundze JB, Mwanza FB, Tshibangu DST, Masengo CA, Liesse L, Takaisi K. Anti-sickling and bacterial inhibitory effects of two medicinal foods from the Congo River basin: *Gnetum africanum* Welw. (Gnetaceae) and *Grewia coriacea* Mast. (Malvaceae). *Current Trad. Med.* (Accepted for publication).
39. Diagne I, Diagne-Gueye ND, Signate-Sy H, Camara B, Lopez-Sall P, Diack-Mbaye A, Sarr M, Ba M, Sow HD, Kuakuvi N. Management of children with sickle cell disease in Africa: experience in a cohort of children at the Royal Albert Hospital in Dakar. *Méd. Trop.* 2003; 63(4-5): 513-520.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>